

Bovine Pinkeye Checklist

Prevent now. Save later.



Risk Factors

Risk Factor:	Prevention Measures:
FACE FLIES	Use dust bags, insecticide ear tags, pour-ons, oral larvicides (IGR's), low-pressure sprayers, mist blower sprayers
WOODY GRASSES	Cut mature long-stem grasses, spray herbicide for woody grasses
EXCESSIVE WETNESS/MANURE	Remove any areas that promote fly breeding ground. Keep dust to a minimum
SAND	Water down sandy/gravel roads near cattle, use masonry sand or washed concrete sand for bedding



Treat now. Save later.

PRODUCT	Presentation	Dose	Booster Dose	Route of Administration	Diarrhoea chemoprophylaxis	CI: suppurative	CI: non-suppurative	CI: suppurative	CI: peritonsillar Type C	CI: peritonsillar Type D	Malignant edema	Malignant edema
Bovilis® Piliguard® Pinkeye+7	10 dose 50 dose	3 mL	3 mL, 3-4 weeks later	SC*	X	X	X	X	X	X	X	X
Bovilis® Piliguard® Pinkeye	10 dose 50 dose	2 mL		SC							X	
Bovilis® 20/20 Vision® 7 with Spur®	10 dose 50 dose	2 mL	3 mL, 3-4 weeks later	SC	X	X	X	X	X	X	X	X
Moraxella bovoculi bacterin	10 dose 50 dose	2 mL	2 mL, 21 days later	SC								X



¹Blackleg

²Malignant edema

³Black disease

⁴Enterotoxemia

⁵Pinkeye, or infectious bovine keratoconjunctivitis

⁶Subcutaneous injection

